

A note on the four-dimensional Kirby calculus

BOGUSŁAW BRODA

ABSTRACT. An explicit derivation of the elements of the representation ring of $SU(2)$ needed to implement the four-dimensional Kirby calculus is sketched.

MAY 1993

1991 *Mathematics Subject Classification*. Primary 57N13; Secondary 57M25, 57N10, 57R65.

Key words and phrases. Four-manifolds, Kauffman bracket, Kirby calculus, surgery presentation, topological invariants.

The aim of this short note is to explicitly derive, in an elementary way, the elements Ω , ω of the representation ring of the group $\text{SU}(2)$ needed to implement the four-dimensional Kirby calculus (handle slides) of César de Sá [1], used in the construction of the surgical invariant of ref. [2].

Sliding an upper curve over a lower one is schematically depicted as

$$\bigcup_{\odot_{\omega}}^{\Omega} \longleftrightarrow \bigcup_{\odot}^{\omega}_{\Omega}, \quad (1)$$

where

$$\Omega = \delta_m W_m, \quad \omega = \bigoplus_{n=0}^N \Delta_n W_n, \quad (2)$$

with W_n an irreducible module of $\text{SU}(2)$, $\dim W_n = n + 1$ (we use the terminology of ref. [3]).

Applying the formula for a connected sum, as well as the satellite formula [3], we obtain, for each m , the fundamental relation [4] (see also [5])

$$\dim_q(W_m) \Delta_n = \sum_k C_{km}^n \Delta_k, \quad n : \dim_q W_n \neq 0, \quad (3)$$

where C_{km}^n are the classical Clebsch-Gordan coefficients, and the quantum dimension

$$\dim_q W_m = (-)^m \frac{q^{\frac{m+1}{2}} - q^{-\frac{m+1}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}. \quad (4)$$

The deformation parameter

$$q = e^{\frac{2\pi i}{r}}, \quad (5)$$

where r is the level, a positive integer.

Thus, we have the following cases:

$$\underline{m=1} \ (r \geq 2)$$

Explicitly, eq. (3) is of the form

$$-\left(q^{\frac{1}{2}} + q^{-\frac{1}{2}}\right) \Delta_n = \Delta_{n+1} + \Delta_{n-1}, \quad n \neq -1 \bmod r, \quad (6)$$

with the initial conditions

$$\Delta_{-1} = 0, \quad \Delta_0 = a_0, \quad (7)$$

a_0 an arbitrary complex number. The only non-trivial solution is given by

$$\Delta_n = \begin{cases} a_0 \dim_q W_n, & \text{if } 0 \leq n \leq r-2; \\ 0, & \text{otherwise.} \end{cases} \quad (8)$$

$m=2$ ($r \geq 3$)

Now, eq. (3) takes the form

$$(q + q^{-1}) \Delta_n = \Delta_{n+2} + \Delta_{n-2}, \quad n \not\equiv -1 \pmod{r}, \quad (9)$$

and decouples for even and odd n . Assuming the initial conditions

$$\Delta_{-2} = \Delta_{-1} = 0, \quad \Delta_0 = a_+, \quad \Delta_1 = a_- \dim_q W_1, \quad (10)$$

where a_+ , a_- are arbitrary complex numbers, we obtain

$$\Delta_n = \begin{cases} a_+ \dim_q W_n, & \text{for } r \text{ odd, if } n \text{ even and } 0 \leq n \leq r-3; \\ a_- \dim_q W_n, & \text{for } r \text{ even, if } n \text{ odd and } 1 \leq n \leq r-3; \\ 0, & \text{otherwise.} \end{cases} \quad (11)$$

$n \geq 3$

There are no non-trivial, i.e. non-zero with a finite N , solutions of eq. (3).

Taking into account that any W_n can be built of W_1 , and any W_n with n even can be built of W_2 , we obtain for r odd the desired moves

$$\begin{array}{ccc} \begin{array}{c} \bigcup_{\omega_0} \\ \odot_{\omega_0} \end{array} \longleftrightarrow \begin{array}{c} \omega_0 \\ \bigcup_{\omega_0} \\ \odot_{\omega_0} \end{array}, & \begin{array}{c} \bigcup_{\omega_+} \\ \odot_{\omega_0} \end{array} \longleftrightarrow \begin{array}{c} \omega_0 \\ \bigcup_{\omega_+} \\ \odot_{\omega_+} \end{array}, & \begin{array}{c} \bigcup_{\omega_+} \\ \odot_{\omega_+} \end{array} \longleftrightarrow \begin{array}{c} \omega_+ \\ \bigcup_{\omega_+} \\ \odot_{\omega_+} \end{array}, \end{array} \quad (12a)$$

but

$$\begin{array}{c} \bigcup_{\omega_0} \\ \odot_{\omega_+} \end{array} \not\longleftrightarrow \begin{array}{c} \omega_+ \\ \bigcup_{\omega_0} \\ \odot_{\omega_0} \end{array}, \quad (12b)$$

where

$$\omega_0 = \bigoplus_{n=0}^{r-2} a_0 \dim_q(W_n) W_n, \quad \omega_+ = \bigoplus_{\substack{n=0 \\ n \text{ even}}}^{r-3} a_+ \dim_q(W_n) W_n. \quad (13)$$

The author is indebted to Prof. H. D. Doebner for his kind hospitality in Clausthal. The work was partially supported by the CEC grant CIPA3510PL921596, the KBN grant 202189101, and the University of Łódź grant.

REFERENCES

1. E. César de Sá, *A link calculus for 4-manifolds*, Topology of low-dimensional manifolds, Proc. Second Sussex Conf., Lecture Notes in Math., vol. **722**, Springer, Berlin, 1979, 16–30.
2. B. Broda, *Surgical invariants of four-manifolds*, E-preprint hep-th/9302092.
3. H. R. Morton and P. Strickland, Math. Proc. Camb. Phil. Soc. **109** (1991) 83–103.
4. B. Broda, *Chern-Simons theory on an arbitrary manifold via surgery*, E-preprint hep-th/9305051.
5. K. Walker, *On Witten's 3-manifold Invariants*, preprint (1990).

DEPARTMENT OF THEORETICAL PHYSICS, UNIVERSITY OF ŁÓDŹ, POMORSKA
149/153, PL-90-236 ŁÓDŹ, POLAND

Current address: Arnold Sommerfeld Institute for Mathematical Physics, Technical University of Clausthal, Leibnizstraße 10, D-W-3392 Clausthal-Zellerfeld, Federal Republic of Germany

E-mail address: ptbb@ibm.rz.tu-clausthal.de